

**VAERS DATA: ALL REPORTED DEATHS OF UNITED STATES RESIDENTS FOLLOWING VACCINATION
6/1/1990 THROUGH 7/23/2021**



Report Run Date: 7/30/2021 (data captured through 7/23/2021)

Vaccine Type	Reported Deaths	Percent (of 10,672 reported events)
ADENOVIRUS TYPE 4 & 7 VACCINE, LIVE ORAL	1	0.01%
ANTHRAX VACCINE	31	0.29%
BACILLUS CALMETTE-GUERIN VACCINE	4	0.04%
COMVAX	193	1.81%
COVID19 VACCINE	6,183	57.94%
DIPHTHERIA AND TETANUS TOXOIDS ACELLULAR PERTUSSIS POLIOVIRUS INACTIVATED HAEMOPHILUS INFLUENZA B AND HEPATITIS B VACCINE (HEXAVAX)	3	0.03%
DIPHTHERIA AND TETANUS TOXOIDS AND ACELLULAR PERTUSSIS VACCINE	693	6.49%
DIPHTHERIA AND TETANUS TOXOIDS AND ACELLULAR PERTUSSIS VACCINE + INACTIVATED POLIOVIRUS VACCINE	8	0.07%
DIPHTHERIA AND TETANUS TOXOIDS AND ACELLULAR PERTUSSIS VACCINE + HEPATITIS B + INACTIVATED POLIOVIRUS VACCINE	406	3.80%
DIPHTHERIA AND TETANUS TOXOIDS AND ACELLULAR PERTUSSIS VACCINE + INACTIVATED POLIOVIRUS VACCINE + HAEMOPHILUS B CONJUGATE VACCINE	222	2.08%
DIPHTHERIA AND TETANUS TOXOIDS AND PERTUSSIS VACCINE	672	6.30%
DIPHTHERIA AND TETANUS TOXOIDS PERTUSSIS AND HAEMOPHILUS INFLUENZA B VACCINE (HEXAVAX)	222	2.08%
DIPHTHERIA AND TETANUS TOXOIDS, PEDIATRIC	12	0.11%
DIPHTHERIA/PERTUSSIS/POLIO (ORAL [LIVE] OR INACTIVATED NOT NOTED)	1	0.01%
DIPHTHERIA/TETANUS/PERTUSSIS/HEPATITIS B	1	0.01%
HAEMOPHILUS B CONJUGATE VACCINE	1,367	12.81%
HAEMOPHILUS B POLYSACCHARIDE VACCINE	9	0.08%
HEPATITIS A	99	0.93%
HEPATITIS A AND HEPATITIS B VACCINE	9	0.08%
HEPATITIS B VACCINE	904	8.47%
HUMAN PAPILLOMAVIRUS (TYPES 6, 11, 16, 18) RECOMBINANT VACCINE	192	1.80%
HUMAN PAPILLOMAVIRUS (TYPES 6, 11, 16, 18, 31, 33, 45, 52, 58) RECOMBINANT VACCINE	26	0.24%
HUMAN PAPILLOMAVIRUS VACCINE	26	0.24%
HUMAN PAPILLOMAVIRUS BIVALENT	5	0.05%
INFLUENZA (H1N1) MONOVALENT (INJECTED)	58	0.54%
INFLUENZA (H1N1) MONOVALENT, (INTRANASAL SPRAY)	9	0.08%
INFLUENZA VIRUS VACCINE, NO BRAND NAME	292	2.74%
INFLUENZA VIRUS VACCINE, QUADRIVALENT (INJECTED)	116	1.09%
INFLUENZA VIRUS VACCINE, QUADRIVALENT (INTRANASAL SPRAY)	10	0.09%
INFLUENZA VIRUS VACCINE, QUADRIVALENT, ADJUVANT (INJECTED)	1	0.01%
INFLUENZA VIRUS VACCINE, QUADRIVALENT, CELL-CULTURE-DERIVED (INJECTED)	7	0.07%
INFLUENZA VIRUS VACCINE, QUADRIVALENT, RECOMBINANT (INJECTED)	4	0.04%
INFLUENZA VIRUS VACCINE, TRIVALENT (INJECTED)	645	6.04%
INFLUENZA VIRUS VACCINE, TRIVALENT (INTRANASAL SPRAY)	21	0.20%
INFLUENZA VIRUS VACCINE, TRIVALENT, ADJUVANT (INJECTED)	8	0.07%
INFLUENZA VIRUS VACCINE, TRIVALENT, CELL-CULTURE-DERIVED (INJECTED)	2	0.02%
INFLUENZA(H1N1) MONOVALENT, UNKNOWN MANUFACTURER	22	0.21%
JAPANESE ENCEPHALITIS VIRUS VACCINE	2	0.02%
JAPANESE ENCEPHALITIS VIRUS VACCINE, INACTIVATED, ADSORBED	1	0.01%
LYME VACCINE (LYMERIX)	9	0.08%
MEASLES VACCINE	8	0.07%
MEASLES, MUMPS AND RUBELLA VIRUS VACCINE, LIVE	254	2.38%
MEASLES, MUMPS, RUBELLA, AND VARICELLA VACCINE (PROQUAD)	24	0.22%
MENINGOCOCCAL B VACCINE	7	0.07%
MENINGOCOCCAL POLYSACCHARIDE VACCINE	40	0.37%
MENINGOCOCCAL VACCINE (MENACTRA)	45	0.42%
MUMPS VIRUS VACCINE, LIVE	4	0.04%
PERTUSSIS, ADSORBED VACCINE	3	0.03%
PLAGUE VACCINE	3	0.03%
PNEUMOCOCCAL VACCINE, POLYVALENT	199	1.86%
PNEUMOCOCCAL, 13-VALENT VACCINE (PREVNAR)	395	3.70%
PNEUMOCOCCAL, 7-VALENT VACCINE (PREVNAR)	687	6.44%
POLIOVIRUS VACCINE INACTIVATED	577	5.41%
POLIOVIRUS VACCINE TRIVALENT, LIVE, ORAL	839	7.86%
RABIES VIRUS VACCINE	16	0.15%
ROTAVIRUS (NO BRAND NAME)	7	0.07%
ROTAVIRUS VACCINE (ROTASHIELD)	10	0.09%
ROTAVIRUS VACCINE, LIVE, ORAL	89	0.83%
ROTAVIRUS VACCINE, LIVE, ORAL, PENTAVALENT	372	3.49%
SMALLPOX VACCINE	17	0.16%
TETANUS AND DIPHTHERIA TOXOIDS AND ACELLULAR PERTUSSIS VACCINE (BOOSTRIX/ADACEL)	67	0.63%
TETANUS AND DIPHTHERIA TOXOIDS, ADULT	31	0.29%
TETANUS TOXOID	10	0.09%
TETRAMUNE	3	0.03%
TYPHOID VACCINE	18	0.17%
VARIVAX-VARICELLA VIRUS LIVE	162	1.52%
YELLOW FEVER VACCINE	24	0.22%
ZOSTER VACCINE	208	1.95%
UNKNOWN VACCINES	169	1.58%
TOTAL INSTANCES REPORTED	16,784	157.27%
TOTAL DEATHS REPORTED	10,672	100.00%

Source: CDC VAERS Database (Vaccine Adverse Events Reporting System)

***NOTE:** There are 10,672 distinct deaths of U.S. residents reported in VAERS through 7/23/2021, many of which occurred after multiple vaccines were administered during the same medical appointment. Therefore there are 16,784 instances of death following vaccinations reported when displayed by vaccine type. But each value displayed in this report includes only distinct patient records (no duplicates) and percentages are calculated as a percent of total patient reports received. For instance, if a patient received influenza, measles, and pertussis vaccines during the same office visit, that patient's record would populate three instances in three categories on this report.