

**VAERS DATA: ALL REPORTED DEATHS OF UNITED STATES RESIDENTS FOLLOWING VACCINATION
6/1/1990 THROUGH 8/27/2021**



Report Run Date: 9/4/2021 (data captured through 8/27/2021)

| Vaccine Type | Reported Deaths | Percent (of 11,382 reported events) |
|---|-----------------|-------------------------------------|
| ADENOVIRUS TYPE 4 & 7 VACCINE, LIVE ORAL | 1 | 0.01% |
| ANTHRAX VACCINE | 31 | 0.27% |
| BACILLUS CALMETTE-GUERIN VACCINE | 4 | 0.04% |
| COMVAX | 193 | 1.70% |
| COVID19 VACCINE | 7,086 | 62.26% |
| DIPHtheria AND TETANUS TOXOIDS ACELLULAR PERTUSSIS POLIOVIRUS INACTIVATED HAEMOPHILUS INFLUENZA B AND HEPATITIS B VACCINE (HEXAVAX) | 3 | 0.03% |
| DIPHtheria AND TETANUS TOXOIDS AND ACELLULAR PERTUSSIS VACCINE | 694 | 6.10% |
| DIPHtheria AND TETANUS TOXOIDS AND ACELLULAR PERTUSSIS VACCINE + INACTIVATED POLIOVIRUS VACCINE | 8 | 0.07% |
| DIPHtheria AND TETANUS TOXOIDS AND ACELLULAR PERTUSSIS VACCINE + HEPATITIS B + INACTIVATED POLIOVIRUS VACCINE | 407 | 3.58% |
| DIPHtheria AND TETANUS TOXOIDS AND ACELLULAR PERTUSSIS VACCINE + INACTIVATED POLIOVIRUS VACCINE + HAEMOPHILUS B CONJUGATE VACCINE | 223 | 1.96% |
| DIPHtheria AND TETANUS TOXOIDS AND PERTUSSIS VACCINE | 672 | 5.90% |
| DIPHtheria AND TETANUS TOXOIDS PERTUSSIS AND HAEMOPHILUS INFLUENZA B VACCINE (HEXAVAX) | 222 | 1.95% |
| DIPHtheria AND TETANUS TOXOIDS, PEDIATRIC | 13 | 0.11% |
| DIPHtheria/PERTUSSIS/POLIO (ORAL [LIVE] OR INACTIVATED NOT NOTED) | 1 | 0.01% |
| DIPHtheria/TETANUS/PERTUSSIS/HEPATITIS B | 1 | 0.01% |
| HAEMOPHILUS B CONJUGATE VACCINE | 1,369 | 12.03% |
| HAEMOPHILUS B POLYSACCHARIDE VACCINE | 9 | 0.08% |
| HEPATITIS A | 100 | 0.88% |
| HEPATITIS A AND HEPATITIS B VACCINE | 9 | 0.08% |
| HEPATITIS B VACCINE | 905 | 7.95% |
| HUMAN PAPILLOMAVIRUS (TYPES 6, 11, 16, 18) RECOMBINANT VACCINE | 193 | 1.70% |
| HUMAN PAPILLOMAVIRUS (TYPES 6, 11, 16, 18, 31, 33, 45, 52, 58) RECOMBINANT VACCINE | 28 | 0.25% |
| HUMAN PAPILLOMAVIRUS VACCINE | 26 | 0.23% |
| HUMAN PAPILLOVAVIRUS BIVALENT | 5 | 0.04% |
| INFLUENZA (H1N1) MONOVALENT (INJECTED) | 58 | 0.51% |
| INFLUENZA (H1N1) MONOVALENT, (INTRANASAL SPRAY) | 9 | 0.08% |
| INFLUENZA VIRUS VACCINE, NO BRAND NAME | 292 | 2.57% |
| INFLUENZA VIRUS VACCINE, QUADRIVALENT (INJECTED) | 117 | 1.03% |
| INFLUENZA VIRUS VACCINE, QUADRIVALENT (INTRANASAL SPRAY) | 10 | 0.09% |
| INFLUENZA VIRUS VACCINE, QUADRIVALENT, ADJUVANT (INJECTED) | 1 | 0.01% |
| INFLUENZA VIRUS VACCINE, QUADRIVALENT, CELL-CULTURE-DERIVED (INJECTED) | 7 | 0.06% |
| INFLUENZA VIRUS VACCINE, QUADRIVALENT, RECOMBINANT (INJECTED) | 4 | 0.04% |
| INFLUENZA VIRUS VACCINE, TRIVALENT (INJECTED) | 645 | 5.67% |
| INFLUENZA VIRUS VACCINE, TRIVALENT (INTRANASAL SPRAY) | 21 | 0.18% |
| INFLUENZA VIRUS VACCINE, TRIVALENT, ADJUVANT (INJECTED) | 8 | 0.07% |
| INFLUENZA VIRUS VACCINE, TRIVALENT, CELL-CULTURE-DERIVED (INJECTED) | 2 | 0.02% |
| INFLUENZA(H1N1) MONOVALENT, UNKNOWN MANUFACTURER | 22 | 0.19% |
| JAPANESE ENCEPHALITIS VIRUS VACCINE | 2 | 0.02% |
| JAPANESE ENCEPHALITIS VIRUS VACCINE, INACTIVATED, ADSORBED | 1 | 0.01% |
| LYME VACCINE (LYMERIX) | 9 | 0.08% |
| MEASLES VACCINE | 8 | 0.07% |
| MEASLES, MUMPS AND RUBELLA VIRUS VACCINE, LIVE | 255 | 2.24% |
| MEASLES, MUMPS, RUBELLA, AND VARICELLA VACCINE (PROQUAD) | 24 | 0.21% |
| MENINGOCOCCAL B VACCINE | 7 | 0.06% |
| MENINGOCOCCAL POLYSACCHARIDE VACCINE | 40 | 0.35% |
| MENINGOCOCCAL VACCINE (MENACTRA) | 45 | 0.40% |
| MUMPS VIRUS VACCINE, LIVE | 4 | 0.04% |
| PERTUSSIS, ADSORBED VACCINE | 3 | 0.03% |
| PLAGUE VACCINE | 3 | 0.03% |
| PNEUMOCOCCAL VACCINE, POLYVALENT | 200 | 1.76% |
| PNEUMOCOCCAL, 13-VALENT VACCINE (PREVNAR) | 396 | 3.48% |
| PNEUMOCOCCAL, 7-VALENT VACCINE (PREVNAR) | 687 | 6.04% |
| POLIOVIRUS VACCINE INACTIVATED | 577 | 5.07% |
| POLIOVIRUS VACCINE TRIVALENT, LIVE, ORAL | 839 | 7.37% |
| RABIES VIRUS VACCINE | 20 | 0.18% |
| ROTAVIRUS (NO BRAND NAME) | 7 | 0.06% |
| ROTAVIRUS VACCINE (ROTASHIELD) | 10 | 0.09% |
| ROTAVIRUS VACCINE, LIVE, ORAL | 89 | 0.78% |
| ROTAVIRUS VACCINE, LIVE, ORAL, PENTAVALENT | 372 | 3.27% |
| SMALLPOX VACCINE | 17 | 0.15% |
| TETANUS AND DIPHtheria TOXOIDS AND ACELLULAR PERTUSSIS VACCINE (BOOSTRIX/ADACEL) | 68 | 0.60% |
| TETANUS AND DIPHtheria TOXOIDS, ADULT | 31 | 0.27% |
| TETANUS TOXOID | 10 | 0.09% |
| TETRAMUNE | 3 | 0.03% |
| TYPHOID VACCINE | 18 | 0.16% |
| VARIVAX-VARICELLA VIRUS LIVE | 163 | 1.43% |
| YELLOW FEVER VACCINE | 24 | 0.21% |
| ZOSTER VACCINE | 212 | 1.86% |
| UNKNOWN VACCINES | 184 | 1.62% |
| TOTAL INSTANCES REPORTED | 17,727 | 155.75% |
| TOTAL DEATHS REPORTED | 11,382 | 100.00% |

Source: CDC VAERS Database (Vaccine Adverse Events Reporting System)

***NOTE:** There are 11,382 distinct deaths of U.S. residents reported in VAERS through 8/27/2021, many of which occurred after multiple vaccines were administered during the same medical appointment. Therefore there are 17,727 instances of death following vaccinations reported when displayed by vaccine type. But each value displayed in this report includes only distinct patient records (no duplicates) and percentages are calculated as a percent of total patient reports received. For instance, if a patient received influenza, measles, and pertussis vaccines during the same office visit, that patient's record would populate three instances in three categories on this report.